



Project ECHO[®] Management of Opioid Use Disorder Clinic

“Suicide Risk Assessment in the Ambulatory Setting”

access this presentation, go to:

<https://razor.med.und.edu/cme/selfserve/web/online-courses>

Presented by

Rory Beil, ReThink Mental Health – Clay County Public Health, Moorhead, MN

Grant Lannoye, PA-C, Towner County Medical Center, Cando, ND

Facilitator: Lisa Faust, MD

Upon completion of this program, the learner will be able to:

- Identify who can administer the C-SSRS
- Identify the Key Biopsychosocial Risk Factors associated with suicide risk
- Reduce false positives, false negatives, and unnecessary interventions

Who should attend?

All members of the healthcare team to include, but not limited to: physicians, NPs, PAs, nurses, behavioral health providers, social workers, pharmacists, medication assisted treatment (MAT) prescribing providers, and prescribing

Why attend? Because you will...

- Participate in case-based clinical discussions focused on the treatment of opioid use disorder.
- Gain up-to-date knowledge of opioid use disorder treatment through brief didactic presentations on topics of interest to learners.
- Improve your confidence in managing patients with opioid use disorder.

For additional information contact: Julie Reiten at 701-777-4173 or julie.a.reiten@und.edu

“The University of North Dakota School of Medicine and Health Sciences is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.”

“The University of North Dakota School of Medicine and Health Sciences designates this enduring activity for a maximum of 1.5 AMA PRA Category 1 Credit(s)[™]. Physicians should claim credit commensurate with the extent of their participation in the activity.”

This project is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under contract number HSH250201600015C. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.